

Frequency and Types of Diabetic Retinopathy in Type II Diabetes; A Hospital Base Study

Mahtab Alam Khanzada, Ashok Kumar Narsani, Faheemullah Shaikh,
Shafi Muhammad Jatoi

ABSTRACT

OBJECTIVES: To determine the frequency and type of diabetic retinopathy in different age groups.

STUDY DESIGN: Descriptive Case series study.

PLACE AND DURATION OF STUDY: Department of Ophthalmology Liaquat University of Medical and Health Sciences (LUMHS) Hyderabad, from February 2009 to January 2010.

METHODS: Two hundred and forty four patients of diabetes mellitus (DM) were randomly selected and grouped out into 30-40 years, 41-50 years, 51-60 years, 61-70 years and more than 70 years of age. Each patient was evaluated for diabetic retinopathy (DR) by fundoscopy and Fundus Fluorescence Angiography (FFA). The retinopathy was graded as 0-3 grade; grade 0= no DR, grade 1= mild DR, grade 2= moderate to severe DR and grade 3= proliferative DR. The different risk factors (age, gender, duration of DM, treatment type and hypertension) were evaluated in relation to diabetic retinopathy.

RESULT: Among 244 patients, 149 were males and 95 were females. Diabetic retinopathy was detected in 100 (40.94%) patients. Mean duration of DM was 13 years in patients with DR and 7.5 years in patients without DR. Most of the DR was found in 40-60 years of age. Out of 244 subjects 25% patients were found with grade 1, 6.96% patients with grade 2 and 9.01% patients were found with grade 3 diabetic retinopathy.

CONCLUSION: Most of the diabetic retinopathy cases were below the age of 60-years, and majority of DR cases presented with type 1 diabetic retinopathy.

KEY WORDS: Diabetic retinopathy, Diabetes Mellitus, Frequency.

INTRODUCTION

In the century 21st with marked improvement in the quality of health care in the various institutes of the under developed countries, the diabetes mellitus has emerged as one of the major health problem worldwide.

Pakistan stands at seven number and by the year 2025 it will be at fifth position in the list of highest diabetic populated country as estimated by International Diabetic Federation (IDF) Database on Diabetics¹. The National Blindness Survey carried out in Pakistan during year 2002-2004 estimated 1.5 million blind peoples and out of these approximately 0.5% became blind due to diabetic complications like diabetic retinopathy (DR)².

Diabetic retinopathy is the most common and devastating microvascular complication of diabetes, which is producing severe visual loss³, and global data also supports this assumption that in the future the diabetic retinopathy will be the leading cause of blindness^{4,5}. The ratio of blindness in type 1 diabetes is 4% and in type 2 it is 1.6%⁶. Previous surveys conducted in Pakistan showed that more than 10% of adult population had type 2 diabetes⁶.

The prevalence of diabetes is reaching epidemic proportion but its frequency varies in different ethnic group⁷. The factors that influence the prevalence of DR include duration of diabetes⁸, types of treatment⁹, hyperglycemia¹⁰, hypertension¹¹, proteinuria¹², serum cholesterol and triglyceride.¹³

The object of this study was to determine the frequency of diabetic retinopathy in type 2 Diabetes mellitus in population of catchment area of LUMHS eye hospital and also to evaluate the status of retinopathy in the study group. This study may help to enhance the awareness of this serious complication of diabetes and to guide the population for early intervention to prevent the further deterioration of vision.

SUBJECTS & METHODS

This descriptive case series study was conducted in the department of Ophthalmology of Liaquat University of Medical and Health Sciences (LUMHS) Hyderabad from February 2009 to January 2010.

Patients of either sex with age of 30 years and above were selected for this study by conventional randomization method. The diagnostic criteria for diabetes mellitus (DM) was set as defined by World Health Or-

ganization (WHO) ¹⁴. Replacement was made for any dropout case attributing to difficulty in grading the retinopathy as a result of concomitant corneal or lenticular opacities, and retinopathy in non-diabetics.

After obtaining the informed consent, information regarding demographics and complete medical history was recorded in a pre-designed proforma including history of treatment regimen, age at onset of diabetes, duration of diabetes and history of hypertension.

A random serum glucose level of each patient was calibrated by single touch gluco- meter to diagnose diabetic control. Seated blood pressure of each patient was measured in the right arm, with mercury sphygmomanometer.

At least two readings of systolic blood pressure (SBP) and diastolic blood pressure (DBP) were taken and mean of two readings was recorded. SBP > 140 mm Hg or DBP > 90 mm Hg was diagnosed as hypertension.

Then pupil of each patient was dilated with 1% tropicamide and 10% phenylephrine to examine the fundus detail, using 90 D with the help of slit lamp binocular microscope. After this fundus fluorescence angiography (FFA) was performed as needed, then grading of severity of diabetic retinopathy was made out as defined by the Early Treatment Diabetic Retinopathy Study (ETDRS) ¹⁵.

Mean and range was calculated for quantitative variables such as age and duration of DM; whereas frequency and percentage were calculated for qualitative variables like gender, types of DR. SPSS 11.0 was used for calculations.

RESULTS

Among 244 subjects 149 were males and 95 were females with mean age of 45±11.5 years. Diabetic retinopathy was present in 100 (49.94%) patients. Duration of DM, type of treatment, gender distribution and presence of hypertension are evaluated in **Table I**. Out of 244 diabetics, 144 did not present DR (grade 0); while among rest of 100 subjects 61 (25.0%) pre-

sented with grade 1 DR, 17 (6.96%) presented with grade 2 DR and 22 (9.0%) presented with grade 3 DR, as detailed in **Table II**.

TABLE I: RISK FACTORS FOR DR (n=244)

Variables	DR (+ve) n=100	DR (-ve) n=144
Duration of DM (Years)		
• Range	5-20	3-16
• Mean	13±4.5	7±3.8
Treatment Regimen		
• Insulin	16 (16.0%)	44 (30.55%)
• Oral	84 (84.0%)	100 (69.44%)
Gender (n, %)		
• Male	51 (51.0%)	44 (30.55%)
• Female	49 (49.0%)	100 (69.44%)
Hypertension	19 (19.0%)	50 (34.72%)

DISCUSSION

In this study DR was present in 40.64% that is approximate to those found in the study from Egypt, that reported the frequency of DR 42.0% and a study from Oman that reported the 42.4% of DR^{9,16}. A previous local study in Pakistan has reported the approximate figure 43.0 % of DR¹⁷. But recently a local study reported 27.43% of DR in the diabetes cases¹⁸. Some studies reported 23.7%, 16.5% and 30.0% DR in India, UK and Spain respectively¹⁹⁻²¹.

The most common type of DR in this study was Grade I which was prevalent in 25.0% of the diabetic patients. These findings are similar to the results of other Asia-based studies^{8,22-25} but the result of grade 2 and 3 of DR showed dissimilar number which contradicted with most of the reports^{13, 17}.

The disparity in our results with recent studies may be due to lack of awareness in our patients to undertake

TABLE II: FREQUENCY AND TYPE OF DR IN DIFFERENT AGE GROUP (n=244)

Age	Grade 0 n (%)	Grade 1 n (%)	Grade 2 n (%)	Grade 3 n (%)	DR +ve n (%)
30-40 year	39 (15.98)	19 (07.78)	03 (01.22)	02 (00.81)	24 (09.83)
41-50 year	15 (06.14)	16 (06.55)	07 (02.86)	03 (01.22)	23 (09.42)
51-60 year	26 (10.65)	16 (06.55)	04 (01.63)	05 (02.04)	25 (10.24)
61-70 year	42 (17.21)	05 (02.04)	03 (01.22)	05 (02.04)	16 (06.55)
71+ year	22 (13.11)	05 (02.04)	00 (00.00)	07 (02.86)	12 (04.91)
Total	144 (59.01)	61 (25.0)	17(06.96)	22 (09.01)	100 (40.94)

regular eye examination or due to the selection bias as only those patients who were already known diabetics having longer duration and who presented with some clinical relevant problems were included in present study.

In our study most of the DR was found between 40-60 years of age. The prevalence of DR in the UAE was 19% and significantly affected elderly males²⁶. In a population-based study of retinopathy from Victoria (the Melbourne Visual Impairment Project [Melbourne VIP]), the frequency of retinopathy was 29.1% in those with self-reported diabetes (age \geq 40 years, types 1 and 2 diabetes)²⁷. In the Blue Mountains Eye Study the frequency of retinopathy was 35.5% based on self-reported diabetes (age \geq 49 years, types 1 and 2 diabetes)²⁸. In present study known diabetics of age \geq 30 years presenting in clinical setting were included among which 40.94% presented with diabetic retinopathy. This difference of age and setting seems to be the reason for difference in results of these studies and the present study.

Duration of diabetes mellitus is directly proportional to the frequency of diabetic retinopathy as proved by Haddad et al⁸ and McKay et al²⁷. Mitchell et al²⁹ reported that about 8% of patients become DR for each year that duration of diabetes increased, it mean as the patient aged with the extension of duration of diabetes, the frequency of DR stepped up and is an independent risk factor for DR.

It was also found in this study that hypertension has no relation to DR, while earlier reports were inconclusive regarding that association between systemic hypertension and DR³⁰. later specific studies found no relation between hypertension and diabetic retinopathy¹⁰.

CONCLUSION

In the present study most of the diabetic retinopathy cases were below the age of 60-years and majority of DR cases presented with type 1 diabetic retinopathy.

REFERENCES

1. International Diabetic Federation Atlas. 2006 showing prevalence of diabetes in 2007 and future projection for 2025 [Internet]. [Updated 2009]. Available from: <http://www.eatlas.idf.org/index1397.html>.
2. Jadoon MZ, Dineen B, Bourne RR, Shah SP, Khan MA, Johnson GJ, et al. Prevalence of blindness and visual impairment in Pakistan: the Pakistan national blindness and visual impairment survey. *Invest Ophthalmol Vis Sci* 2006; 47:4749-55.
3. Mensah E, Kohner E. Diagnosis and management of diabetic retinopathy. *Topic Endocrinol* 2002;19:14-18.
4. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes. *Diabetic Care* 2004; 27:1047-53.
5. American Diabetic Association. Diabetic Retinopathy. *Diabetes Care* 2002; 25:S90-3.
6. Shera AS, Rafique G, Khwaja IA, Baqai S, Khan IA, King H. Pakistan National Diabetes Survey prevalence of glucose intolerance and associated factors in North West at Frontier Province (NWFP) of Pakistan. *J Pak Med Assoc* 1999; 49:206-11.
7. Gharaagaji R, Fagihzadeh S, Meshkani M R, Rohipor R. The Prevalence of Diabetic Retinopathy and Related Risk Factors in Diabetic Type I, in Tehran Iran *Res J Biol Sci* 2008; 3 (6): 596-600.
8. EL Haddad OA, Saad MK. Prevalence and risk factors for diabetic retinopathy among Omani diabetics. *Br J Ophthalmol* 1998;82:901-6.
9. Santos K.G, Tschiedel B, Schneider JR, Souto KEP, Roisenberg I. Prevalence of retinopathy in Caucasian type 2 diabetic patients from the South of Brazil and relationship with clinical and metabolic factors. *Braz J Med Biol Res*, February 2005; 38(2): 221-25.
10. Van Leiden HA, Dekker JM, Moll AC, Nijpels G, Heine RJ, Bouter LM. Blood pressure, lipids, and obesity are associated with retinopathy. The Hoorn Study. *Diabetes Care* 2003; 25: 1320-1325.
11. Tapp RJ, Shaw JE, Harper CA, de Courten MP, Balkau B, McCarty DJ. The prevalence of and factors associated with diabetic retinopathy in Australian population. *Diabetes Care* 2003; 26 [26]:1731-7.
12. The EURODIAB IDDM Complications Study. Retinopathy and vision loss on insulin-dependent diabetes in Europe. *Ophthalmology* 1997; 104:252-60.
13. Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and it's complications: estimates and projections to the year 2010. *Diabet Med* 1997 14(suppl.5):551-85.
14. World Health Organization. The WHO diagnostic criteria of Diabetes of 1999 as revised in 2006 at the 19th World Diabetes Conference [Internet]. [updated 2009]. Available from: <http://www.who.int/Diabetes/publications/en>.
15. Early Treatment Diabetic Retinopathy Study (ETDRS) Research Group. Grading diabetic retinopathy from stereoscopic color fundus photographs-an extension of the Modified Airlie House Classification. ETDRS Report Number 10. *Ophthalmology* 1991;98(Suppl):786-806.
16. Herman WH, Aubert RE, Engelgau MM, et al. Diabetes mellitus in Egypt glycaemic control and microvascular and neuropathic complications. *Diabet Med* 1998;15:1045-51.

17. Shera AS, Jawad F, Maqsood A, Jamal S, Azfar M, Ahmed U. Prevalence of chronic complication and associated factors in Type 2 Diabetes. *J Pak Med Assoc* Feb 2004; 54(2):54-9.
18. Mahar PS, Awan MZ, Manzar N, Memon MS. Prevalence of type-II diabetes mellitus and diabetic retinopathy: The Gaddap Study. *J College of Physicians Surgeons Pak* 2010; 20(8): 528-32.
19. Ramachandaran A, Snehalatha C, Satyavani K. Prevalence of complications and their risk factors in type 2 diabetes. *J Assoc Physicians India* 1999; 47:1152-6.
20. Morgen CL, Currie CJ, Hunt J. The prevalence of multiple diabetes related complications. *Diabet Med* 2000; 17 :146-51.
21. Zafra Mexcua JA, Mendez Segovia JC, Novalbos Ruiz JP. Chronic complications in patients with type 2 diabetes mellitus cared for at a health center. *Aten Primaria* 2000 ;25:529-35.
22. Jamal-u-Din A, Qureshi MB, Khan AJ, Khan MD, Ahmad K. Prevalence of diabetic retinopathy among individuals screened positive for diabetes in five community-based eye camps innorthern Karachi, Pakistan. *J Ayub Med Coll Abbottabad* 2006; 18:41-3.
23. Wong TY, Cheung N, Tay WT, Wang JJ, Aung T, Saw SM, et al. Prevalence and risk factors for diabetic retinopathy: the Singapore Malay eye study. *Ophthalmology* 2008; 115:1869-75.
24. Khan AJ. Age sex and duration relationship and prevalence of Diabetic retinopathy in Pakistani population. *Pak J Ophthalmol* 1990; 6:6-8.
25. Rani PK, Raman R, Chandrakantan A, Pal SS, Perumal GM, Sharma T. Risk factors for diabetic retinopathy in self-reported rural population with Diabetes. *J Postgrad Med* 2009; 55:92-6.
26. Al-Maskari F, El-Sadig M. Prevalence of diabetic retinopathy in the United Arab Emirates community-based eye camps in northern Karachi, Pakistan. *J Ayub Med Coll Abbottabad* 2006; 18:41-3.
27. McKay R, McCarty C A, Talyor H R. Diabetic retinopathy in Victoria, Australia: the Visual Impairment Project. *Br J Ophthalmol* 2000; 84:865-70.
28. Mitchell P, Smith W, Wang JJ, Attebo K. Prevalence of diabetic retinopathy in an older community. The Blue Mountains Eye Study *Ophthalmology* 1998; 105:406–11: a cross-sectional Survey. *BMC Ophthalmology* 2007, 7:11 <http://www.biomedcentral.com/1471-2415/7/11/prepub>
29. Mitchell P, Moffitt P. Update and implications from the Newcastle diabetic retinopathy study *Aust N Z J Ophthalmol* 1990;18:13-17.
30. Klein R, Klein BEK, Moss SE. The Wisconsin Epidemiologic Study of Diabetic Retinopathy III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol* 1984; 102:527–32.



AUTHOR AFFILIATION:

Dr. Mahtab Alam Khanzada (Corresponding Author)
Assistant Professor, Department of Ophthalmology
Liaquat University of Medical and Health Sciences
(LUMHS), Jamshoro/Hyderabad, Sindh-Pakistan.
E-mail: dr.khanzada@yahoo.com

Dr. Ashok kumar Narsani
Associate Professor, Department of Ophthalmology
LUMHS, Jamshoro/Hyderabad, Sindh-Pakistan.

Dr. Faheemullah Shaikh
Assistant Professor, Department of Ophthalmology
LUMHS, Jamshoro/Hyderabad, Sindh-Pakistan.

Dr. Shafi Muhammad Jatoi
Professor, Department of Ophthalmology
LUMHS, Jamshoro/Hyderabad, Sindh-Pakistan.