

The Effect of Combination of Citronella Oil and Back Massage on Sleep Quality and Stress Levels in Postpartum Mothers

Heyrani Heyrani^{1*}, Melania Asi¹, Elyasari Elyasari¹, Maya Astuti², Arsulfa Arsulfa¹, Hesty Resyana¹

ABSTRACT

OBJECTIVE: This study aimed to investigate the impact of combining citronella oil with back massage on sleep quality and stress levels in postpartum mothers.

METHODOLOGY: This study employed a quantitative, quasi-experimental design with two groups, utilizing pre- and post-tests. This study was conducted from August 2024 to April 2025 in the Wawolesea, Lembo, and Saka sub-districts of the North Konawe Regency. A total of 196 samples were collected using purposive sampling. The Pittsburgh Sleep Quality Index (PSQI) was used to measure sleep quality, and the Perceived Stress Scale-10 was used to measure stress levels. The data were analysed using both univariate and bivariate methods.

RESULTS: The intervention group experienced 79 cases of negative sleep quality, with a p-value of 0.000, indicating that the intervention significantly improved sleep quality. In contrast, the control group showed seven positive cases with a p-value of 0.157, which was not statistically significant. For stress levels, the intervention group reported 86 negative cases, with a p-value of 0.003, indicating significant effectiveness, whereas the control group reported a p-value of 0.076. The intervention group had a mean sleep quality score of 24.50 compared to 36.50 in the control group ($p < 0.001$), as well as a stress level score of 25.47 versus 35.53 ($p = 0.011$).

CONCLUSION: The combination of *citronella oil* and back massage not only improves sleep quality but also significantly reduces stress levels in postpartum mothers.

KEYWORDS: *Citronella Oil*, Back Massage, Sleep Quality, Stress Level, Postpartum Mother

INTRODUCTION

Physiological, emotional, and psychological changes in postpartum mothers are indicators of the postpartum phase. These changes may affect mothers' general health and well-being. Postpartum mothers frequently experience increased stress and poor sleep quality. Non-pharmacological methods, such as massage therapy and the use of citronella oil, can be employed to reduce stress and improve sleep quality. The plant that yields citronella oil, *Cymbopogon nardus*, is known to promote blood circulation and relaxation¹.

The integration of holistic health practices in postpartum care through a multidisciplinary approach is essential for optimizing maternal well-being. Holistic strategies encompass emotional support alongside traditional medical care, recognized as vital in fostering psychological well-being and enhancing breastfeeding outcomes. A study indicates that emotional release practices, such as self-talk, are

significantly associated with improved maternal mental health and lactation, highlighting the importance of addressing emotional well-being in postpartum care.

The postpartum period is marked by significant physiological and psychological transitions influenced by a combination of biological, psychological, and social factors, as described in the biopsychosocial model. Physiologically, women undergo changes in their reproductive and endocrine systems, as well as alterations in blood circulation and vital signs, reflecting the body's adaptation following delivery. For example, uterine involution and hormonal adjustments are crucial for recovery, highlighting the biological aspect of postpartum adaptations³.

The integration of massage and aromatherapy into postpartum care routines offers significant benefits for new mothers, particularly in alleviating physical and emotional stressors associated with childbirth. Studies indicate that techniques such as endorphin massage can effectively reduce anxiety levels in postpartum mothers, enhancing their psychological well-being⁴. Combining this with loving massage and aromatherapy, particularly using lavender oil, has been shown to improve breast milk production and promote relaxation, thereby supporting maternal recovery⁵.

Massage therapy and *citronella oil* exert calming effects through complex physiological and neurobiological mechanisms. Massage therapy is recognized for its ability to stimulate pressure

¹Department of Midwifery, Faculty of Medicine, Polytechnic of the Ministry of Health Kendari, Kendari, Southeast Sulawesi, Indonesia

²Department of Midwifery, Faculty of Medicine, Polytechnic of the Ministry of Health Bandung, Bandung, West Java, Indonesia

Correspondence: heyрани1112@gmail.com

doi: 10.22442/jlumhs.2025.01306

Received: 28-04-2025

Revised: 30-06-2025

Accepted: 11-07-2025

Published Online: 29-07-2025



receptors in the skin, which can activate the vagus nerve, leading to increased vagal activity associated with reduced cortisol levels related to stress⁶. This reduction in cortisol has been linked to improved mood and relaxation, as evidenced in various populations, including infants and the elderly⁷. Additionally, massage promotes an increase in serotonin and dopamine, neurotransmitters that play crucial roles in mood regulation and contribute to feelings of well-being⁸. *Citronella oil*, an essential oil known for its calming properties, enhances this relaxing effect. Its aromatic compounds are believed to influence neurobiological responses through olfactory pathways, which can affect the limbic system, particularly in emotional regulation and stress response⁹. The combined application of massage therapy with aromatherapy using citronella may

The postpartum phase involves complex physiological, emotional, and psychological transitions that significantly impact the mother's health and well-being. Postpartum depression (PPD) rates can be as high as 21%, highlighting the urgent need for effective intervention strategies¹⁰. New mothers often experience increased stress and decreased sleep quality, which can exacerbate health problems, including anxiety and hypertension¹¹. Nonpharmacological interventions, such as massage therapy combined with aromatherapy, have shown potential in addressing these challenges. For example, studies have shown that aromatherapy with essential oils, including those with calming properties, can improve relaxation and emotional well-being among postpartum mothers¹¹. Additionally, the efficacy of combining massage with aromatherapy to enhance relaxation and reduce muscle tension supports better sleep quality in this population¹². These dual interventions can help create an environment that promotes maternal recovery¹³.

This study aimed to examine the effects of *citronella oil* and back massage on postpartum stress levels and sleep quality. The novelty of this study lies in the fact that numerous studies have been conducted on this topic; however, this study offers a fresh perspective by investigating the combined effects of massage techniques and the use of *citronella oil*. By utilising massage techniques in conjunction with citronella oil, we aim to enhance our understanding of how to improve the welfare of postpartum mothers and provide an additional resource for complementary postpartum care methods.

METHODOLOGY

A two-group pre-test–post-test design with a quasi-experimental approach was employed in this quantitative study. Questionnaires and observational techniques were used to gather data. From August 2024 to April 2025, the study was conducted in the Wawolesea, Lembo, and Saka sub-districts of the North Konawe Regency.

This study included all postpartum women in the North

Konawe Regency in 2024. The Lemeshow formula was used to calculate the sample size, and 196 mothers were included in the sample. The sampling technique employed was purposive. The inclusion criteria for the included samples were as follows: age between 17 and 45 years, experiencing sleep issues or stress as measured before the intervention, being in the postpartum period (0–6 weeks postpartum), and being willing to provide informed consent. The exclusion criteria were serious health issues, history of severe mental disorders, inability to adhere to the intervention process thoroughly, and use of sedatives were the exclusion criteria.

Mothers were approached through community health centers and local maternal health clinics within the Wawolesea, Lembo, and Saka sub-districts. The recruitment process included disseminating information about the study objectives and eligibility criteria in various community forums and through healthcare providers. Interested participants were screened for inclusion and exclusion criteria, ensuring a focused cohort of mothers who could benefit from the intervention involving *citronella oil*.

In this study, the fidelity of the intervention involving *citronella oil* was supported by comprehensive training provided to the nurses and midwives participating in the fieldwork. The training encompassed specific objectives, procedural protocols, and operational definitions related to the intervention. It included sessions on the therapeutic properties and application methods for *citronella oil*, emphasizing the importance of consistency in dosage and administration techniques to maintain the integrity of the integration process. Moreover, the training addressed best practices in participant engagement and ethical considerations, promoting informed consent and participant safety. An emphasis was placed on the accurate use of assessment tools, including the Pittsburgh Sleep Quality Index (PSQI) and the Perceived Stress Scale-10, to ensure that participants received a standardized approach to monitoring outcomes. This preparatory phase aimed to cultivate competency and confidence among nurses and midwives, thereby establishing a robust foundation for the reliability and validity of the intervention.

Applying for a research permit was the first step of this study, followed by site exploration. *Citronella oil* was used as an intervention material. The primary researcher enlisted the assistance of qualified nurses and midwives to support the fieldwork. After that, socialisation sessions were conducted to help all the research assistants in the field understand the goals and direction of the study. Stress levels and sleep quality of postpartum mothers were assessed through a screening process. Informed consent was obtained, and samples were split into intervention and control groups, which were then gathered according to the inclusion criteria. Following socialization, the intervention was conducted as planned. After data processing and creation of an extensive report, the

evaluation was completed with a post-test to gauge the research outcomes.

The research timeline began in August 2024, with the application for a research permit and site exploration, as well as the initiation of recruitment of mothers through community health centres and clinics. September 2024: Pre-tests conducted, assessing sleep quality (using the Pittsburgh Sleep Quality Index, PSQI) and stress levels (using the Perceived Stress Scale-10) among the participants who met the inclusion criteria. October 2024: Training for nurses and midwives on intervention protocols, including objectives and application of *citronella oil*, concluded. November 2024: Intervention commenced, where the intervention group received *citronella oil* while the control group did not.

February 2025: Post-tests conducted to evaluate changes in stress and sleep quality in both groups. April 2025: Data analysis finalized, along with the preparation of results for dissemination.

As the study design incorporated two independent groups (intervention and control), bivariate analysis was employed to assess differences in outcomes between these groups. The Wilcoxon test, a non-parametric test, was appropriately selected because the data on stress levels and sleep quality do not exhibit a normal distribution. This choice enables reliable comparisons of ordinal or continuous data that do not meet the assumptions of parametric tests. Furthermore, the Mann-Whitney U test was used to examine post-treatment variations in stress levels and sleep quality between the two groups, thereby reinforcing the validity of the findings when normality assumptions are untenable. The choice of these tests thus ensured robust statistical inference while addressing pertinent research questions regarding the efficacy of *citronella oil* as an intervention.

Sleep quality was evaluated by using the Pittsburgh Sleep Quality Index (PSQI). This is due to the high validity and reliability of the Pittsburgh Sleep Quality Index (PSQI), which is used to assess various aspects of sleep. Stress levels were measured using the Perceived Stress Scale-10. The Ethics Commission of the Kendari Ministry of Health Polytechnic granted ethical approval (DP.04.03/F.XXXVI.39/003/2024).

RESULTS

Table I shows the distribution of respondents between the intervention group and the control group in a study. The analysis includes demographic variables such as age, education, occupation, number of live births, sleep quality (both pre-test and post-test), and stress levels (both pre-test and post-test). In a specific period, the intervention group tended to be younger than the control group. Education showed a fairly balanced proportion, although there were slightly more respondents with junior high school education in the control group. In terms of occupation, most respondents were household workers, with relatively low poverty levels in both groups. Sleep quality

improved in the intervention group post-intervention, as indicated by a decrease in the number of respondents reporting disturbed sleep. Stress levels also changed, with more respondents in the intervention group reporting mild stress levels after the intervention compared to the control group. All variables showed good homogeneity values ($p > 0.05$), indicating no significant differences between the two groups in the measured demographic characteristics.

Table I: Distribution of respondents in the intervention and control groups

Variables	Intervention group		Intervention group		Homogeneity sample
	Frequency	%	Frequency	%	
Age					
17-34	69	70.4	76	77.6	0.882
35-44	23	23.5	17	17.3	
>45	6	6.1	5	5.1	
Education					
Elementary school	13	13.3	7	7.1	0.831
Junior high school	7	7.1	20	20.4	
Senior high school	63	64.3	59	60.2	
University	15	15.3	12	12.2	
Work					
Government employee	3	3.1	3	3.1	0.952
Household	88	89.8	81	82.7	
Private employee	5	5.1	7	7.1	
Unemployment	2	2.0	7	7.1	
Number of live births					
Primipara	33	33.7	40	40.8	0.873
Multipara	65	66.3	58	59.2	
Sleep quality (pre-test)					
Not disturbed	35	35.7	26	26.5	0.845
Disturbed	63	64.3	72	73.5	
Sleep quality (post-test)					
Not disturbed	83	84.7	73	74.5	0.817
Disturbed	15	15.3	25	25.5	
Stress level (pre-test)					
Mild	20	20.4	30	30.6	0.811
Moderate	58	59.2	56	57.1	
Heavy	20	20.4	12	12.2	
Stress level (Post test)					
Mild	60	61.2	40	40.8	0.821
Moderate	30	30.6	50	51.0	
Heavy	8	8.2	8	8.2	
Levene test					

As shown in **Table II**, out of 98 participants, the intervention group, which received back massage and citronella oil, experienced 79 cases of poor sleep quality and 0 positive cases. In contrast, the control group experienced only 7 positive cases. The control group recorded 91 cases in the tie category compared to 19 cases in the intervention group. A p-value of 0.000 was found for sleep quality in the analysis, indicating that the intervention significantly improved sleep quality. Furthermore, the intervention group recorded 86 negative cases of stress levels, whereas the control group recorded only 13. The intervention had an impact on stress levels, as evidenced by the p-value for stress levels in the intervention group (0.003).

According to **Table III**, the mean score for sleep quality in the intervention group was 24.50, while that in the control group was 36.50, with a p-value of 0.001. These findings suggest that the two groups differ significantly in the degree to which the intervention improves sleep quality. With a p-value of 0.011, the mean stress level in the intervention group was 25.47, and that in the control group was 35.53. These findings also revealed significant differences between the two groups.

Table II: Differences in sleep quality and stress levels in each group

Cases	Sleep quality		Stress levels	
	Intervention	Control	Intervention	Control
Negative	79	0	86	0
Positive	0	7	7	13
Ties	19	91	6	85
p-value	0.000	0.157	0.003	0.076

Wilcoxon test

Table III: Differences in sleep quality and stress levels after treatment between the two groups

Variables	N	Mean	p-value
Sleep quality			
After treatment			
Intervention	98	24.50	0.001
Control	98	36.50	
Stress levels			
After treatment			
Intervention	98	25.47	0.011
Control	98	35.53	

Mann-Whitney U Test

DISCUSSION

The results indicated that the stress levels and sleep quality of the intervention group significantly changed following treatment. The proportion of individuals with undisturbed sleep quality has increased from 35.7% to 84.7 %. This result is consistent with earlier

research demonstrating that structured intervention programs have a significant impact on stress and sleep quality¹⁴. Furthermore, the results indicated that the percentage of people who experienced extreme stress levels decreased from 20.4% to 8.2% following the intervention. This demonstrates that the intervention can help individuals sleep better and experience less stress. Additionally, prior research has shown that massage can improve sleep quality¹⁵. The results also demonstrated that the intervention and control groups differed significantly from each other. The intervention group reported 79 cases of poor sleep quality, with no positive cases, whereas the control group reported only seven positive cases. Numerous studies have demonstrated that aromatherapy using essential oils, such as peppermint and lavender, can enhance sleep quality, which helps explain this phenomenon^{16,17}. According to earlier research by Chen et al. (2015)¹⁸, the sleep quality of postpartum women can be enhanced by lavender aromatherapy. The results of this study demonstrated effects comparable to those of earlier studies, in which *citronella oil* and massage techniques were combined.

Aromatherapy, particularly with lavender oil, has been shown to influence cortisol levels and sleep patterns. The administration of lavender aromatherapy can stimulate the olfactory system, activating brain regions that regulate emotional responses and stress¹⁹. Research indicates that exposure to lavender essential oil can lead to a reduction in cortisol levels, as observed in various studies measuring cortisol levels in saliva and urine²⁰. One study highlighted a decrease in salivary cortisol levels of 0.8 µg/dL after inhalation, suggesting a calming effect on the sympathetic nervous system²⁰. Furthermore, aromatherapy with lavender has been linked to improved sleep quality by promoting relaxation and reducing anxiety²¹.

In contrast to the control group, which had only 13 negative cases for stress levels, the intervention group had 86 negative cases. Before and after the intervention, the results revealed a p-value of 0.003 in the intervention group. This suggests that the intervention groups differed significantly before and after treatment. There was no difference between the control group before and after treatment, as indicated by the control group's p-value of 0.076, which was lower than that of the intervention group. It is well known that there is a higher chance of stress and depression during the postpartum phase²². According to earlier studies, using specific essential oil types can lessen anxiety and depression^{23,24}. According to earlier studies, lavender oil used as aromatherapy can help haemodialysis patients feel less uncomfortable and improve their overall wellbeing²⁴.

By lowering stress, sleep quality was improved in the intervention group. This result implies a connection between the two elements. Stress reduction has been

linked to better sleep quality, according to earlier research²⁵. Postpartum women's physical and mental health benefit of this positive cycle. Health professionals can use the success of the intervention as a guide when creating comparable programs to enhance sleep quality in postpartum mothers.

Interestingly, in contrast to the intervention group, the control group showed a significantly higher number of positive cases, including stress reduction and improved sleep quality. This serves as the foundation for supporting the idea that massage and *citronella oil* can help postpartum mothers sleep better and feel less stressed. This is corroborated by research by Park et al.²⁶ which shows aromatherapy massage greatly enhances sleep quality and lessens fatigue.

According to the results, the average sleep quality in the intervention group was 24.50, with a p-value = 0.001. In contrast to the control group, which had an average of only 36.50, this indicates that the treatment had a significantly positive effect on sleep quality. As with the stress level variable, the control group's average was 35.53, while the intervention group's average was 25.47 with a p-value of 0.011.

These results are consistent with those of earlier studies, which have demonstrated that the use of essential oils can dramatically lower stress levels and enhance sleep quality. This was evident in a study by Karadağ et al.. The results demonstrated that aromatherapy had a positive impact on patients' anxiety levels and sleep quality, which is consistent with the information gathered in this study²⁷.

When used in conjunction with massage, essential oils can influence biological processes associated with emotions and stress. The nervous system can be influenced by essential oils, which help lower the stress-related hormone cortisol²⁸. This is further supported by Yang et al.'s research, which shows that inhaling lavender oil can reduce stress and enhance sleep quality by regulating the body's hormonal system.¹⁷

In particular, both factors underscore the potential of aromatherapy as a non-invasive approach to enhance mental and physical well-being. According to the study's observations, the intervention's impact aligns with findings from related research, indicating that aromatherapy can provide emotional solace, as evidenced by improved sleep and reduced stress. For instance, Lee et al.'s study demonstrated that aromatherapy can help patients undergoing medical procedures sleep better and experience less pain, suggesting that aromatherapy and relaxation have a significant combined effect³⁰.

The use of aromatherapy and massage as interventions for postpartum women demonstrates potential long-term benefits in enhancing maternal well-being and reducing psychological distress. Aromatherapy has shown efficacy in decreasing postpartum depression (PPD) symptoms through trials that reported lower Edinburgh Postnatal Depression Scale (EPDS) scores in women receiving

aromatherapy compared to control groups^{31,32}. Additionally, routines incorporating aromatherapy along with oxytocin massage have been linked to increased breast milk production and improved emotional states among postpartum mothers^{33,34}.

Cultural perceptions significantly influence the acceptance and effectiveness of aromatherapy and massage therapy across diverse populations. The historical development of these practices varies, as evident in the differences between countries. For instance, the development of aromatherapy in Poland has been less pronounced compared to Western nations, where such modalities are more ingrained in health practices³⁵. Positive attitudes toward these therapies, shaped by cultural beliefs and personal experiences, can either enhance or hinder their perceived efficacy³⁶. Furthermore, individuals' socio-cultural backgrounds, including their spirituality and religious inclinations, influence their openness to complementary and alternative medicine (CAM) practices, which can lead to varying levels of acceptance among groups³⁷. Such insights underscore the importance of culturally sensitive approaches in implementing aromatherapy and massage within public health strategies.

Our study is critical because it may offer medical professionals new approaches to enhancing the general health of new mothers. Additionally, these favourable results may inspire further studies on the integration of massage and aromatherapy in maternal care procedures. We hope that your readers will find this article interesting, as we believe it offers essential perspectives on maternal health and the use of complementary therapies. We hope for a positive review and look forward to hearing from you.

CONCLUSION

Postpartum mothers who received interventions that combined back massage with *citronella oil* reported significantly improved sleep and lower stress levels. Overall, it can be concluded that the intervention was suitable for treating sleep issues, although the analysis revealed that the intervention group had lower quality sleep. The control group, on the other hand, did not address stress levels but demonstrated improved sleep quality. The results indicated that the intervention had a significant impact on both measured variables. The effectiveness of this approach in providing postpartum emotional and physical support, which can impact overall well-being, was demonstrated by the intervention group's notable reduction in stress levels. These results suggest that to promote the health and recovery of postpartum mothers, clinical practice should consider complementary approaches, such as those employed in this study. In this regard, back massage and the application of *citronella oil* can be regarded as potentially beneficial methods for enhancing the mental health and quality of life of new mothers. Further investigation is required to examine the

variations in individual responses and confirm the long-term efficacy of these interventions.

Ethical permission: Kendari Ministry of Health Polytechnic, Ethics committee, Indonesia, ERC letter No. DP.04.03/F.XXXVI.39/003/2024.

Conflict of Interest: The author states no conflict of interest.

Financial Disclosure/Grant Approval: Funding grant from the Ministry of Health of Indonesia with Number HK.02.03/F.XXXVI/1666/2024.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

Heyrani H: Conception, Interpretation, Analysis, Drafting, Revising, Final approval

Asi M: Interpretation, Drafting, Revising, Final approval

Elyasari E: Interpretation, Drafting, Revising, Final approval

Astuti M: Revising, Final Approval

Arsulfa A: Revising, Final Approval

Resyana H: Revising, Final Approval

REFERENCES

1. Chrisna A, Cahyati Y, Mardiani D. Effectiveness Of Giving Ginger And Citronella Oil For Dysmenorrhea In Adolescent Girls. *Jordan J Nurs Res.* 2024;2(3):135–42.
2. Haloho CBR, Urnia EE, Astuti DR, Murti NN. Enhancing Breastfeeding Self-Efficacy In The Postpartum Period Through Emotional Release With Self-Talk Method. *Healthc Low Resour Settings.* 2024 Jan 30;11(1):1–11.
3. Martin C, Viky M, Rokibullah R, Sofinia H. The Physiological Changes In The Postpartum Period After Childbirth. *Asian J Soc Human.* 2022 Dec 12;1(03):105–18.
4. Widyawati M, Nazilah R, Latifah L. Mezile's Model (Back Massage And Self-Hypnosis) Based On Information And Technology To Lose The Anxiety Of Postpartum Mothers In Covid-19 Pandemic Era. *Open Access Maced J Med Sci.* 2022 Jan 3;10(T8):188–93.
5. Widiastuti NMR, Arini KN, Yuniati MG. Midwifery Complementary Treatment With The Application Of Oxytocin Massage Using Lavender Aromatherapy Oil On Breast Milk Production In Postpartum Mothers. *Babali Nurs Res.* 2022 Nov 30;3(3):246–53.
6. Cole JS, Olson AD, Dupont-Versteegden EE. The Effects Of Massage Therapy In Decreasing Pain And Anxiety In Post-Surgical Patients With Breast Cancer: A Systematic Review And Meta-Analysis. *Glob Adv Integr Med Health.* 2024 Jan 16;13(1):1–11.
7. Widyaningrum R, Ulu THM, Yulianti AD. A

Literature Review: Effectiveness Of Swedish Foot Massage Therapy On Quality Of Sleep In Elderly With Essential Hypertension. *Dis Prevent Public Health J.* 2023 Jun 27;17(2):134–40.

8. Bingöl H, Akaras E, Kocaman H. The Effects Of Massage Therapy On Symptoms Related Cerebral Palsy In Children With Cerebral Palsy: A Systematic Review. *J Clin Med Kazakhstan.* 2020 Aug 31;4(58):6–14.
9. Jastrzębska E, Górecka-Bruzda A, Ogłuszka M, Lipka Ms, Pawłowska A. Effect Of Massage On Stress Indicators In Recreational Horses—A Pilot Study. *Animals.* 2025 Mar 11;15(6):789–99.
10. Yamakawa Y, Maruta M, Higuchi Y, Tokunaga A, Iwanaga R, Honda S et al. Factors Influencing Postpartum Depression Among Japanese Parents: A Prospective Longitudinal Study. *Neuropsychopharmacol Rep.* 2023 Jun 13;43(2):213–21.
11. Mor S, Sela Y, Lev-Ari S. Postpartum Mothers' Mental Health In A Conflict-Affected Region: A Cross-Sectional Study Of Emotion Regulation And Social Support. *J Clin Med.* 2025 Feb 13;14(4):1244–54.
12. Singh DR, Sunuwar DR, Adhikari S, Singh S, Karki K. Determining Factors For The Prevalence Of Depressive Symptoms Among Postpartum Mothers In Lowland Region In Southern Nepal. *PloS One.* 2021 Jan 22;16(1):1–11.
13. Walker LO, Murry N. Maternal Stressors And Coping Strategies During The Extended Postpartum Period: A Retrospective Analysis With Contemporary Implications. *Women's Health Reports.* 2022 Jan 1;3(1):104–14.
14. Nourian M, Nikfarid L, Khavari Am, Barati M, Allahgholipour Ar. The Impact Of An Online Mindfulness-Based Stress Reduction Program On Sleep Quality Of Nurses Working In Covid-19 Care Units. *Holist Nurs Pract.* 2021;35(5):257–63.
15. Habibi Nezhad M, Sadeghi Nk, Yaghoubinia F. Comparison Of The Effect Of Swedish Massage And Earplugs/Eye Masks On Sleep Quality Of Patients Admitted To ICUs. *Med Surg Nurs J.* 2024; 12(2): 1–8.
16. Cheng H, Lin L, Wang S, Zhang Y, Liu T, Yuan Y et al. Aromatherapy With Single Essential Oils Can Significantly Improve The Sleep Quality Of Cancer Patients: A Meta-Analysis. *BMC Complement Med Ther.* 2022;22(1):1–12.
17. Yang CP, Yang CC, Tsai IJ, Lin TH, Chiou YI, Wang HF et al. The Immediate Effects Of Lavender-Based Essential Oil Inhalation On Subsequent Polysomnography In People With Poor Sleep Quality. *J Chinese Med Assoc.* 2023;86(7):665–71.
18. Chen S, Chen C. Effects Of Lavender Tea On Fatigue, Depression, And Maternal-Infant Attachment In Sleep-Disturbed Postnatal Women. *Worldviews Evid Based Nurs.* 2015;12(6):370–9.
19. Sani FN, Irdianty MS. The Effects Of Slow Stroke

- Back Massage And Lavender Aromatherapy On Blood Pressure In Hypertensive Patients. *Indonesian J Med*. 2020;5(3):178–84.
20. Can Çiçek S, Demir Ş, Yılmaz D, Açıkgoz A, Yıldız S, Yis ÖM. The Effect Of Aromatherapy On Blood Pressure And Stress Responses By Inhalation And Foot Massage In Patients With Essential Hypertension. *Holist Nurs Pract*. 2022 Jul;36(4):209–22.
 21. Ogata K, Ataka K, Suzuki H, Yagi T, Okawa A, Fukumoto T et al. Lavender Oil Reduces Depressive Mood In Healthy Individuals And Enhances The Activity Of Single Oxytocin Neurons Of The Hypothalamus Isolated From Mice: A Preliminary Study. *Evid Based Complement Altern Med*. 2020 Jan 14;2020(1):1–12.
 22. Luberto CM, Park ER, Goodman JH. Postpartum Outcomes And Formal Mindfulness Practice In Mindfulness-Based Cognitive Therapy For Perinatal Women. *Mindfulness (N Y)*. 2018;9(3):850–9.
 23. Chen MI, Chen YE, Lee HF. The Effect Of Bergamot Essential Oil Aromatherapy On Improving Depressive Mood And Sleep Quality In Postpartum Women: A Randomized Controlled Trial. *J Nurs Res*. 2022;30(2):1–8.
 24. Vafaei S, Shorofi A, Kamali M, Asgarirad H, Mousavinasab SN, Azadbakht M et al. The Effect Of Topical Application Of Lavender Lipogel On Sleep Quality And Well-Being In Patients Undergoing Maintenance Hemodialysis: A Randomized Clinical Trial. *Jundishapur J Chronic Dis Care*. 2023;12(4):1–12.
 25. Rahmani N, Araj-Khodaei M, Mohammad-Alizadeh-Charandabi S, Esmaeilpour K, Montazeri M, Mirghafourvand M. The Effect Of Cognitive–Behavioral Counseling With Or Without Citrus Aurantium Essential Oil On Sleep Quality In Pregnant Women: A Randomized Controlled Trial. *Sleep Biol Rhythms*. 2023;21(3):337–46.
 26. Park H, Chun Y, Kwak S. The Effects Of Aroma Hand Massage On Fatigue And Sleeping Among Hospice Patients. *Open J Nurs*. 2016;06(07):515–23.
 27. Karadag E, Samancioglu S, Ozden D, Bakir E. Effects Of Aromatherapy On Sleep Quality And Anxiety Of Patients. *Nurs Crit Care*. 2017;22(2):105–12.
 28. Sattayakhom A, Wichit S, Koomhin P. The Effects Of Essential Oils On The Nervous System: A Scoping Review. *Molecules*. 2023;28(9):1–23.
 29. Kongkaew C, Sakunrag I, Chaiyakunapruk N, Tawatsin A. Effectiveness Of Citronella Preparations In Preventing Mosquito Bites: Systematic Review Of Controlled Laboratory Experimental Studies. *Tropical Medicine & International Health*. 2011;16(7):802–10.
 30. Lee J, Hur MH. The Effects Of Aroma Essential Oil Inhalation On Stress, Pain, And Sleep Quality In Laparoscopic Cholecystectomy Patients: A Randomized Controlled Trial. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2022;16(1):1–8.
 31. Nguyen NT, Pengpid S. Proactive Approaches To Preventing Postpartum Depression In Non-Depressive Pregnant Women: A Comprehensive Scoping Review. *Front Glob Womens Health*. 2025;6(4):1–12.
 32. Hu T, Lee S, Loh E. Effectiveness Of Aromatherapy For Intrapartum And Postpartum Emotional Problems Among Parturient Women: A Meta-Analysis Of Randomized Controlled Trials. *Japan J Nurs Sci*. 2022 Jul 2;19(3):1–11.
 33. Widiastuti NMR, Arini KN, Yuniati MG. Midwifery Complementary Treatment With The Application Of Oxytocin Massage Using Lavender Aromatherapy Oil On Breast Milk Production In Postpartum Mothers. *Babali Nurs Res*. 2022 Nov 30;3(3):246–53.
 34. Sahriana S, Jasmawati J, Nulhakim L. Enhancement Of Breast Milk Production Through The Combination Of Oxytocin Massage And Lavender Aromatherapy In Postpartum Mothers. *Int J Nurs Midwif Sci*. 2023 Nov 3;7(2a):64–71.
 35. Hartman-Petrycka M, Lebedowska A. The Assessment Of Quality Of Products Called Sandalwood Oil Based On The Information Provided By Manufacturer Of The Oil On Polish, German, And English Websites. *Evidence-Based Complement Altern Med*. 2021 Jul 12;2021(7):1–8.
 36. Dehghan M, Samareh Fekri A, Rashidipour N, Naeimi Bafghi N, Maghfouri A, Ebadzadeh M. The Effects Of Aromatherapy With Clove And Lavender On Headache Caused By Spinal Anesthesia In Patients Undergoing Urological Surgery: A Randomized Clinical Trial Study. *Health Sci Rep*. 2025 Feb 25;8(2):1–11.
 37. Akova I, Ayilgan AP, Mercan C, Duman EN, Adigüzeloğlu Dağdeviren M, Dolmuş Dirican Mg. Medical School Students' Attitudes, Behaviours, And Information Sources Towards Holistic Complementary And Alternative Medicine: A Cross-Sectional Study. *J Traditional Med Complement Ther*. 2024;7(1):23–32.

