

Enhancing HIV/AIDS Prevention in Adolescents: An Information- Motivation-Behavioral Skills Approach

Muhtar Muhtar^{1*}, Julhana Julhana¹, Sri Djuwitaningsih², Komang Ayu Henny Achjar³, Fadlurrahmi Fadlurrahmi¹

ABSTRACT

OBJECTIVE: To determine how adolescents' knowledge and attitudes regarding HIV/AIDS prevention were affected by health education using a motivational information approach.

METHODOLOGY: This study employed a quasi-experimental design conducted in Bima, NTB, Indonesia, from July 2024 to January 2025. A cluster sampling technique was used, yielding a total of 374 students divided into two groups. The control group received HIV/AIDS educational materials in the form of textbooks without support from cadres. In contrast, the intervention group was given a booklet about HIV/AIDS and received guidance from a cadre. The intervention was carried out four times a week for six months. Data analysis was performed using a paired t-test.

RESULTS: The intervention significantly improved knowledge and attitudes towards HIV/AIDS prevention in adolescents, regardless of the control or intervention group. The intervention group showed a significant increase in knowledge scores about HIV/AIDS, with an average score of 13.74, compared to the control group's 8.04. The intervention group also showed a significant increase in attitude scores, with an average of 47.12, compared to the control group's average of 35.04. Both knowledge and attitude in the intervention group had p-values of <0.000.

CONCLUSION: The intervention proved to be more effective in improving knowledge and attitudes towards HIV/AIDS.

KEYWORDS: HIV-AIDS, Health Education, Cadre, Information, Adolescents.

INTRODUCTION

Adolescents are the fastest growing population and are at risk for HIV/AIDS in all countries.¹ AIDS transmission currently mainly attacks teenagers of various age groups.² Nearly 50% of adolescents diagnosed with HIV are under the age of twenty-five. HIV is currently the leading cause of death for individuals between the ages of 15 and 24.³ This happens because they tend to use condoms incorrectly, and teenagers are generally infected through unprotected sexual activity and drug use as a result of promiscuity.⁴ Adolescents represent a demographic significantly affected by HIV/AIDS, particularly due to their heightened vulnerability during this transitional life stage. Research indicates that adolescents exhibit risky sexual behaviors, partly driven by misconceptions about HIV transmission and the stigma surrounding the disease.^{5,6} Peer influence and lack of comprehensive sexual education compound this issue, as many youths remain

unaware of accurate prevention methods.⁷

The success of HIV prevention in adolescents depends on the understanding of HIV prevention and transmission among them.⁸ HIV/AIDS prevention strategies can include initiatives, AIDS counseling services, contact centers, and needle exchange programs. Several activities targeting educational interventions play a significant role in raising awareness and encouraging HIV AIDS prevention among adolescents.⁹ Several strategic preventive steps have been taken in the field of education and increasing awareness, especially for teenagers (mainly in the areas of safe sex promotion, risk behaviour, and disease transmission).¹⁰ One of the most important efforts for HIV prevention is encouraging responsibility for actions, using condoms, and promoting healthy behavior.¹¹

A good understanding will indirectly increase awareness of HIV/AIDS, both in terms of behavior and knowledge, so that this will affect the reduction of the risk of infection.¹² Generally, HIV/AIDS is influenced by risky sexual behavior, such as changing partners and unsafe sex behavior, both in terms of condom use and transmission methods.¹³ Adolescent girls have less knowledge of HIV/AIDS than adolescent boys. So adolescent girls often have greater misunderstandings regarding HIV/AIDS than adolescent boys. This affects perceptions of how to prevent this disease.¹⁴ A Study has shown the need for comprehensive sex education to improve knowledge and attitudes regarding risky behaviors for HIV/AIDS transmission.¹⁵

¹Department of Nursing, Poltekkes Kemenkes Mataram, City of Mataram, Province of NTB, Indonesia

²Department of Nursing, Poltekkes Kemenkes Jakarta III, City of Bekasi, province of West Java, Indonesia

³Department of Nursing, Poltekkes Kemenkes Denpasar, City of Denpasar, Province of Bali, Indonesia

Correspondence: muhtarbima998@gmail.com

doi: 10.22442/jlumhs.2025.01272

Received: 21-02-2025

Revised: 14-07-2025

Accepted: 21-07-2025

Published Online: 30-07-2025



Unplanned pregnancy, high-risk behavior, and education programs are essential to improving understanding and attitudes related to HIV/AIDS.¹⁶ Domestic violence, dating behavior, healthy relationships, and social and emotional learning can be improved with comprehensive sex education.¹⁷

Both psychological and socioeconomic factors can significantly influence adolescents' high-risk behaviors regarding sexual activity. Psychological distress, such as anxiety and mood disorders, is closely associated with risky sexual behaviors, including early sexual debut and inconsistent condom use. Furthermore, adolescents facing adverse psychological effects are likely to engage in sexually risky activities as a coping mechanism to deal with feelings such as loneliness or anxiety.¹⁸

The purpose of this study was to analyze the impact of health education with a motivational information approach on adolescents' knowledge and attitudes regarding HIV/AIDS prevention. The study is categorized as new because it provides insight into the effectiveness of this strategy to improve adolescents' understanding and perspectives regarding HIV/AIDS.

METHODOLOGY

Research Design

The design of this study is quasi-experimental, with pre- and post-tests, and compares the control and intervention groups.

Study Area

This study was conducted in Bima City, NTB Province, Indonesia, between July 2024 and January 2025. Bima City has 39 schools, with 5,562 students (data from the Ministry of Education, Indonesia). The sample consisted of high school students, regardless of ethnicity, age, and gender. Cluster sampling was employed in this study due to its practicality and efficiency in the context of Bima City, where the population is distributed across multiple schools. Cluster sampling allows researchers to divide the population into distinct subgroups or clusters (in this case, schools), from which a sample of high school students can be randomly selected. This method is particularly advantageous when dealing with large populations, as it reduces time and costs associated with data collection by enabling researchers to focus on specific locations—in this instance, selected schools within Bima City.

Additionally, cluster sampling mitigates the difficulty of obtaining a comprehensive list of high school students, thereby enhancing feasibility. It ensures the sample remains representative of the school's diverse demographics without requiring extensive data on each student. Ultimately, using cluster sampling in this study aligns with both methodological rigor and logistical practicality, ensuring a sufficiently diverse and representative sample of high school students.

Sampling

The cluster sampling technique is used. With this

technique, the researcher divided the research area into five sub-districts based on Bima City. The formula used is $n = n / (1 + n e^2)$. The number of samples obtained was 374 students divided into 5, so that each specific area received a sample proportion of 74 to 75.

Intervention

The intervention to enhance knowledge and attitudes towards HIV/AIDS comprised a systematic five-stage approach. Initially, communication was established with key local agencies, including the Director General of the Ministry of Education and Culture in Bima, to ensure compliance with existing regulations (Stage One). Subsequently, a baseline survey assessed the knowledge and attitudes of the target population regarding HIV/AIDS, informing the development of a tailored education program (Stage Two).

In the third stage, mentors were organized into five groups strategically positioned across various schools, tasked with providing essential guidance to student samples regarding HIV/AIDS. The program's design differentiated between a control group, which received a textbook module without mentorship, and an intervention group that received both a booklet and mentor support, with structured sessions held four times a week for six months (Stage Four).

Finally, in the last month of the intervention, comprehensive monitoring and evaluation were conducted to measure changes in knowledge and attitudes, comparing results from the initial survey to those following the educational intervention (Stage Five). This rigorous framework ensures that the program is not only compliant with policy but also effective in achieving its academic goals.

Data Collection Tools and Techniques

The questionnaire used to assess the level of knowledge is based on the WHO. To evaluate attitudes, a Likert-scale questionnaire containing 10 questions was used. (1) Strongly disagree, (2) disagree, (3) agree, (4) Strongly agree. The questionnaire used to assess knowledge levels was based on sources from the World Health Organization (WHO) to ensure the accuracy and relevance of the collected information. It was designed to evaluate respondents' understanding of specific health issues addressed by the WHO, such as disease prevention, health promotion, or safe health practices. To assess attitudes, the questionnaire consisted of 10 Likert-scale questions. In this format, respondents were asked to rate specific statements regarding their attitudes and perceptions by selecting one of four options: "strongly disagree" (1), "disagree" (2), "agree" (3), and "strongly agree" (4). The use of a Likert scale allows for measuring the intensity of attitudes, providing an in-depth quantitative analysis of respondents' views on relevant issues. Thus, this questionnaire provides a comprehensive tool for exploring two crucial dimensions—knowledge and attitudes—that influence individual health behaviors.

Data analysis

Data on age, gender, knowledge and attitude were tested univariately. A paired T-test was used to assess the differences in the level of knowledge and attitude between the control and treatment groups. The use of the paired t-test in this study has several underlying reasons. First, the paired t-test is designed to compare two sets of interrelated measurements, in this case, the participants' knowledge and attitudes before and after an intervention (control and treatment groups). This allows researchers to measure changes in the same variables in the same individuals, thereby increasing the accuracy of the results. Second, the paired t-test accounts for variability in individual measurements, thereby reducing type I errors that might occur if the analyses were conducted independently.

Ethical Clearance

This research has received approval from the ethics committee of Poltekkes Kemenkes Mataram with No. 0140/EA/KEPK/2024.

RESULTS

The average age of the adolescents in the control group was 15.72 years, with a standard deviation of 0.991, as seen in **Table I**. Participants in the intervention group were aged 14-18 years. The mean age of the respondents was 16.18 years, accompanied by a standard deviation of 1.101.

Table II presents a comparative analysis of respondent characteristics by gender in two groups: Intervention and Control. The sample consisted of 187 respondents in the Intervention group, with 45% males (n=83) and 55% females (n=104). In the Control group, there were also 187 respondents, with 47.5% males (n=89) and 52.5% females (n=98).

Table III revealed a notable disparity in knowledge levels between the intervention and control groups. The intervention group achieved an average

knowledge score of 8.02, whereas the control group recorded a slightly higher average score of 8.04. Results from the paired T-test demonstrated a significant difference in knowledge levels pre- and post-intervention. Furthermore, the study identified significant variations in attitudes between the two groups. The intervention group had a mean attitude score of 35.04, compared to the control group's mean score of 34.96. The paired T-test results indicated a significant difference in attitudes in the intervention group, whereas the control group showed no significant change.

Table IV indicates that the intervention group showed greater improvement in knowledge than the control group across the four targeted areas of knowledge. These areas include the transmission of HIV, the causative agent of AIDS, the heightened risk associated with multiple sexual partners, and the understanding of HIV carriers.

Table V indicates that the intervention group exhibited a more favourable attitude than the control

Table I: Respondent characteristics based on age

Variables	Mean	SD	Min Max	95% CI
Age				
Control	15.72	0.991	14-18	15.44 – 16.00
Intervention	16.18	1.101	14-18	15.87 - 16.49

Table II: Respondent characteristics based on gender

Variables	Group			
	Intervention		Control	
	n	%	n	%
Gender				
Man	83	45	89	47.5
Woman	104	55	98	52.5

Table III: Knowledge and attitudes before and after treatment in the intervention and control groups in adolescents

Variables	Group	Mean	SD	S.E.	Min Mak	95% CI	P value
Knowledge	Intervention						0.000
	Pre	8.02	1,059	0.150	6 - 10	7.72 – 8.32	
	Post	13.74	0.828	0.117	12 - 15	13.50 – 13.98	
	Control						0.124
	Pre	7.94	0.867	0.123	5 - 10	7.69 - 8.19	
	Post	8.04	0.781	0.111	6 - 10	7.82 – 8.26	
Attitude	Intervention						0.000
	Pre	35.04	1,761	0.237	30 - 40	34.54 – 35.54	
	Post	47.12	1,637	0.237	44 - 52	46.65 – 47.59	
	Control						0.144
	Pre	34.96	1,678	0.248	31 - 40	34.48 - 35.44	
	Post	35.04	1,678	0.332	31 – 40	34.56 – 35.52	

group, as evidenced by the greater percentage increase in attitude scores for both groups. The evaluation of attitude encompassed two key aspects: youth vulnerability to HIV and the willingness to undergo voluntary HIV testing.

Table IV: Knowledge about issues related to HIV/AIDS before and after intervention

Knowledge	Correct answers (Pre)	Correct answers (Post)	Correct answers (Pre)	Correct answers (Post)
	Intervention		Control	
Spread of HIV	77.7 %	81.5 %	76.9 %	79.5 %
Causative agent of AIDS	75.5 %	83.2 %	73%	76.8 %
Multiple sex partners increase the risk	47.5 %	83.5 %	44.5 %	78.5 %
Concept of HIV carrier	12.7 %	23.8 %	17.5 %	20.1 %

Table V: Attitudes about issues related to HIV/AIDS before and after the intervention

Attitude	Attitude (Pre)	Attitude (Post)	Attitude (Pre)	Attitude (Post)
	Intervention		Control	
Youth vulnerability to HIV	70%	83%	66.9 %	79.7 %
HIV voluntary testing	84.3 %	91.5 %	81.5 %	85.5 %

DISCUSSION

The findings showed that the intervention group showed a greater increase in knowledge than the control group. The average score in the intervention group reached 13.74, while in the control group it was only 8.04. This shows that the program in the intervention group was more successful than that in the control group. These results are supported by previous studies, which show that individual understanding increases with effective, targeted educational strategies.¹⁹

This study itself focuses on HIV transmission and the virus that causes HIV/AIDS, by focusing on the knowledge that having multiple sexual partners can be a significant source of risk for HIV/AIDS transmission. This study also highlights the concept of HIV carriers, where individuals who are carriers of the virus are very likely to have no symptoms. The results of the study showed that the intervention group was more effective in answering questions than the control group. This proves that proper education can improve understanding of the dangers of HIV/AIDS.²⁰

HIV is generally transmitted through various ways, including from mother to child during childbirth, use of

unsafe injection needles and unsafe sex. Previous studies have shown that knowledge about this can reduce the risk of HIV/AIDS transmission.²¹ Adequate knowledge will also be the basis for individuals to make decisions regarding dangerous sexual behavior and unsafe use of needles.²² The importance of good knowledge is that individuals can know how the virus is transmitted so that they can avoid risky behavior.²³ With increased education and proper education, communities can avoid the spread of HIV/AIDS. Educating yourself and others about HIV/AIDS prevention is essential to protecting yourself from HIV/AIDS infection. We can reduce the spread of the virus and help people living with HIV by raising awareness and understanding about the disease. A key to combating this is by tackling the spread of HIV.²⁴ proactive steps can contribute to suppressing the spread of HIV/AIDS.

Acquired immunodeficiency syndrome (AIDS) is the result of the human immunodeficiency virus (HIV) compromising the immune system by primarily targeting CD4 T cells.²⁵ Sharing needles, engaging in sexual activity, nursing, or giving birth can all transfer HIV. Nearly every respondent in the intervention group demonstrated a solid grasp of the information covered in the HIV/AIDS education session and provided thoughtful responses. The results show that the intervention approach effectively raises knowledge and awareness of the illness.

Other findings suggest that education about early detection and adherence to therapy can be a lifesaver for individuals with HIV/AIDS.²⁶ Improving the welfare of people affected by HIV/AIDS is the key to success in preventing this disease from spreading further. This is driven by awareness of the establishment of comprehensive prevention and treatment initiatives in cases of people with HIV/AIDS.²⁷

Having several sexual partners increases the risk of contracting HIV because of the higher chance of contracting the virus during unprotected intercourse. The increased likelihood of transmission explains this phenomenon due to the larger number of possible encounters with people who may be HIV-positive. Unprotected sexual activity has been repeatedly demonstrated to be a major contributor to the spread of HIV in studies, highlighting the significance of safe sex practices in reducing this risk.²⁸ To prevent HIV from spreading to others, it's critical to engage in safe sexual behaviour and get tested for the virus regularly. There were no discernible differences between the intervention and control groups' answers on this question. The percentages in the control and intervention groups, which did not change substantially, made this clear. It is vital to take precautions and seek medical advice when at risk because HIV infection can have significant repercussions for both the infected person and their partner.

Every individual needs to receive appropriate information regarding the dangers of having multiple

partners, which can increase the risk of HIV/AIDS transmission. Safe sex practices will generally protect the community from various infectious diseases that may exist. Safe sex practices also minimize the spread of HIV/AIDS.²⁹ Eliminating stigma and increasing awareness of safe sexual behavior can be a pioneer in preventing HIV/AIDS or other sexually transmitted diseases.

People with HIV/AIDS who do not show any symptoms and signs of the disease are often referred to as HIV carriers. People with these characteristics should receive good care by carrying out a routine medication schedule to reduce the risk of transmission to others.³⁰ In both groups, based on this study, they tend to provide a deep response. The intervention given to the intervention group proved effective in increasing knowledge and awareness of HIV/AIDS and the importance of proper treatment.

For those living with HIV, regular evaluation of their viral load and CD4 count is essential to confirming the effectiveness of their treatment regimen. Frequent monitoring enables medical professionals to guarantee patients' optimal health outcomes and make well-informed decisions about therapy modifications.³¹ Support from medical professionals and neighbourhood resources can also assist people in taking their medications as prescribed and preserving their general health. To boost their immune system and overall health, HIV carriers should adopt healthy lifestyle habits, such as regular exercise and a balanced diet. Additionally, counselling programs and mental health support can help people living with HIV deal with any emotional difficulties that may come up.³²

The findings showed that the improvement in attitudes in the intervention group was greater than that in the control group. The intervention group showed an average attitude score of 47.12, compared to the control group's 35.04. This certainly highlights how vital the right intervention program is to influence participants' attitudes towards HIV/AIDS. In addition, a focused education program is significant in understanding the stigma associated with HIV/AIDS.

HIV-AIDS education initiatives have the potential to significantly alter social perceptions and practices, which will ultimately help achieve the overarching objective of eliminating the stigma and discrimination related to the illness. To create a more accepting and inclusive environment for those impacted by HIV/AIDS, policymakers and medical professionals should highlight these efforts. We can promote a more understanding and caring society that is better prepared to handle the difficulties experienced by individuals impacted by HIV/AIDS by funding extensive educational initiatives³³. To stop the disease from spreading, these programs can also support early detection, access to treatment, and preventive measures.

Several factors, including the stigma attached to HIV

and the lack of comprehensive sexual education, can influence the attitudes of young people at risk for the virus.³⁴ Communities must provide the support and tools necessary to empower youth to make informed decisions regarding their sexual health. This can entail encouraging candid discussions about sex, making condoms and other types of protection available, and providing testing and treatment services. By addressing these barriers, we can improve community health overall and reduce the risk of HIV transmission among youth. Compared to the control group, the intervention group, which has been receiving treatment for six months, has a superior understanding of this.

Promoting an inclusive and supportive culture can foster an atmosphere in which adolescents feel safer making informed decisions about their sexual health³⁵. Protecting themselves from HIV infection requires Collaboration among stakeholders to ensure that adolescents are equipped with the resources and knowledge. Educating young people about sexual health and accessible health services is needed. Collaboration between educational institutions, medical facilities, neighborhood associations, and government agencies can make this happen. Together, these parties can develop a comprehensive program.

Depending on personal experiences, anxieties, and beliefs, attitudes regarding voluntary HIV testing can differ significantly.³⁶ While some people may feel nervous or stigmatized, others may see testing as a proactive step in managing their health. To encourage more people to get tested, healthcare providers should provide information and assistance. People may feel more at ease obtaining tests and managing their health if a secure, accepting environment is provided.³⁷ Furthermore, having access to factual information on HIV might aid in debunking misconceptions and lessening the stigma associated with the virus.

Community outreach to address misconceptions and promote the importance of knowing one's HIV status is also needed.³⁸ Healthcare providers address barriers to testing and provide resources for those who may be hesitant. This can help them maintain their crucial role in raising HIV prevention levels and promoting comprehensive public health. Collaboration between government organizations, healthcare providers, and the general population is necessary to create comprehensive HIV prevention and treatment plans.³⁹

LIMITATION

Because this study primarily involved high school students from a single school in the city of BIMA, variations in educational background and socioeconomic conditions across regions may not be reflected.

CONCLUSION

The educational program that was carried out effectively increased understanding of HIV/AIDS. This

is evidence that the intervention carried out raised awareness of HIV/AIDS.

Increased knowledge and attitudes can be seen from various factors such as agents that cause HIV/AIDS, having multiple partners will pose a high risk, and how HIV/AIDS transmission can occur. In general, the intervention group outperformed the control group in various aspects. Overall, the intervention group had a strong understanding of the educational material provided. This is evidence that the intervention was successful in increasing awareness and understanding of this disease. In light of the findings from this study, future researchers are encouraged to expand the scope of interventions aimed at raising HIV/AIDS awareness by incorporating diverse educational strategies, such as interactive workshops or peer-led discussions, which may further enhance engagement and retention of information.

Ethical Permission: Komite Etik Penelitian Kesehatan, Indonesia, ERC letter No. 0140/KEPK/2024.

Conflict of interest: There is no conflict of interest between the authors.

Financial Disclosure / Grant Approval: No funding agency was involved in this research.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION: All authors contribute equally

REFERENCES

1. Eslami Aa, Ebrahimi Z, Rahimi M, Fathian-Dastjerdi Z, Bagherikholejani F. Family-Based Interventions In Youth To Prevent HIV/AIDS: A Systematic Review. *J Educ Health Promot*. 2023;12(1):1–9.
2. Archary M, Pettifor AE, Toska E. Adolescents And Young People At The Centre: Global Perspectives And Approaches To Transform HIV Testing, Treatment And Care. *J Int Aids Soc*. 2020;23(S5):1–4.
3. Shannon CI, Klausner JD. The Growing Epidemic Of Sexually Transmitted Infections In Adolescents: A Neglected Population. *Curr Opin Pediatr*. 2018;30(1):137–43.
4. Murali V, Jayaraman S. Substance Use Disorders And Sexually Transmitted Infections: A Public Health Perspective. *Bjpsych Adv*. 2018;24(3):161–6.
5. Maris Bakara S, Sahara Lubis E, Fitriani Y. Adolescent Knowledge And Perception Of HIV/AIDS Stigmatization In The Indonesian Context. *Jurnal Berkala Epidemiologi*. 2023 Sep 15;11(3):287–94.
6. Siregar PA. Analysis AIDS Stigmatization Of Adolescents In Medan City. *Saintika Medika*. 2021 Jun 10;17(1):61–70.
7. Agu IC, Mbachu CO, Okeke C, Eze I, Agu C, Ezenwaka U et al. Misconceptions About Transmission, Symptoms And Prevention Of HIV/AIDS among Adolescents In Ebonyi State, South-East Nigeria. *BMC Res Notes*. 2020 Dec 14;13(1):1–5.
8. Ratnawati D, Setiawan A, Sahar J, Widyatuti, Nursasi Ay, Siregar T. Improving Adolescents' Hiv/Aids Prevention Behavior: A Phenomenological Study Of The Experience Of Planning Generation Program (Genre) Ambassadors As Peer Educators. *Belitung Nurs J*. 2024;10(1):56–66.
9. Macounová P, Tomášková H, Šnajdrová A, Stanovská M, Polochová M, Tomášek I, Et Al. Education Of Adolescents In The Prevention Of HIV/AIDS In The Czech Republic. *Int J Environ Res Public Health*. 2021;18(11):1–12.
10. Moreira Mt, Rocha E, Lima A, Pereira L, Rodrigues S, Fernandes Cs. Knowledge About Sex Education In Adolescence: A Cross-Sectional Study. *Adolescents*. 2023;3(3):431–45.
11. Santos MJ De O, Ferreira EMS, Ferreira MC. Predictors Of Condom Use Among College Students. *Int J Environ Res Public Health*. 2024;21(4):433.
12. Obeagu EI, Obeagu GU, Ede MO, Odo EO, Buhari HA. Translation Of HIV/AIDS Knowledge Into Behavior Change Among Secondary School Adolescents In Uganda: A Review. *Medicine*. 2023;102(49):1–4.
13. Ren Z, Zhou Y, Liu Y. Factors Associated With Unsafe Sexual Behavior Among Sexually Active Chinese University Students, Hebei Province, 2019. *BMC Public Health*. 2021;21(1):1904.
14. Kawuki J, Gatasi G, Sserwanja Q, Mukunya D, Musaba MW. Comprehensive Knowledge About HIV/AIDS And Associated Factors Among Adolescent Girls In Rwanda: A Nationwide Cross-Sectional Study. *Bmc Infect Dis*. 2023 Jun 7;23(1):382.
15. George G, Beckett S, Reddy T, Govender K, Cawood C, Khanyile D et al. Role Of Schooling And Comprehensive Sexuality Education In Reducing HIV And Pregnancy Among Adolescents In South Africa. *J Acquired Immune Deficiency Syndromes*. 2022 Jul 1;90(3):270–5.
16. Mbizvo MT, Kasonda K, Muntalima NC, Rosen JG, Inambwae S, Namukonda ES et al. Comprehensive Sexuality Education Linked To Sexual And Reproductive Health Services Reduces Early And Unintended Pregnancies Among In-School Adolescent Girls In Zambia. *BMC Public Health*. 2023;23(1):348.
17. Todesco M, Breman J, Haryanto NN, Kok G, Massar K. Effect Evaluation Of A Comprehensive Sexuality Education Intervention Based On Socio-Emotional Learning Among Adolescents In Jakarta, Indonesia. *Front Public Health*. 2023 Oct 2;11.

18. Pengpid S, Peltzer K. Psychological Distress And Its Associated Factors Among School-Going Adolescents In Tanzania. *Psychol Stud (Mysore)*. 2020 Jun 28;65(2):174–81.
19. Raghupathi V, Raghupathi W. The Influence Of Education On Health: An Empirical Assessment Of OECD Countries For The Period 1995–2015. *Archives Of Public Health*. 2020 Dec 6;78(1):20.
20. Alhasawi A, Grover Sb, Sadek A, Ashoor I, Alkhabbaz I, Almasri S. Assessing HIV/AIDS Knowledge, Awareness, And Attitudes Among Senior High School Students In Kuwait. *Medical Principles And Practice*. 2019;28(5):470–6.
21. Yang D, Allen J, Mahumane A, Riddell J, Yu H. Knowledge, Stigma, And HIV Testing: An Analysis Of A Widespread Hiv/Aids Program. *J Dev Econ*. 2023;160:102958.
22. Kassie AA, Gudayu TW, Araya BM. Knowledge, Attitude, And Preventive Practices Towards Sexually Transmitted Infections Among Preparatory School Students In West Gojjam Zone, Ethiopia. *Adv Public Health*. 2020 May 12;2020:1–9.
23. Pavlopoulou Id, Dikaloti Sk, Gountas I, Sypsa V, Malliori M, Pantavou K et al. High-Risk Behaviors And Their Association With Awareness Of Hiv Status Among Participants Of A Large-Scale Prevention Intervention In Athens, Greece. *BMC Public Health*. 2020 Dec 28;20(1):105.
24. Bouabida K, Chaves BG, Anane E. Challenges And Barriers To HIV Care Engagement And Care Cascade: Viewpoint. *Frontiers In Reproductive Health*. 2023 Jul 20;5.
25. Masenga Sk, Mweene Bc, Luwaya E, Muchaili L, Chona M, Kirabo A. HIV Host Cell Interactions. *Cells*. 2023;12(10):1351.
26. Jocelyn, Nasution FM, Nasution NA, Asshiddiqi MH, Kimura NH, Siburian MHT et al. HIV/AIDS in Indonesia: Current Treatment Landscape, Future Therapeutic Horizons, And Herbal Approaches. *Front Public Health*. 2024;12.
27. Gandhi RT, Bedimo R, Hoy JF, Landovitz RJ, Smith DM, Eaton EF et al. Antiretroviral Drugs For Treatment And Prevention Of HIV Infection In Adults. *JAMA*. 2023;329(1):63.
28. Du X, Zhang L, Luo H, Rong W, Meng X, Yu H et al. Factors Associated With Risk Sexual Behaviours Of HIV/STDS Infection Among University Students In Henan, China: A Cross-Sectional Study. *Reprod Health*. 2021;18(1):1–11.
29. Wirawan GBS, Gustina NIZ, Januraga PP. Open Communication About Reproductive Health Is Associated With Comprehensive Hiv Knowledge And A Non-Stigmatizing Attitude Among Indonesian Youth: A Cross-Sectional Study. *Journal Of Preventive Medicine And Public Health*. 2022 Jul 31;55(4):342–50.
30. Seyedalinaghi S, Mirzapour P, Pashaei Z, Afzalian A, Tantuoyir MM, Salmani R et al. The Impacts Of Covid-19 Pandemic On Service Delivery And Treatment Outcomes In People Living With Hiv: A Systematic Review. *Aids Res Ther*. 2023 Jan 6;20(1):4.
31. Shoko C, Chikobvu D. A Superiority Of Viral Load Over CD4 Cell Count When Predicting Mortality In Hiv Patients On Therapy. *BMC Infect Dis*. 2019 Dec 15;19(1):169.
32. Fauk NK, Merry MS, Mwanri L, Hawke K, Ward PR. Mental Health Challenges And The Associated Factors In Women Living With HIV Who Have Children Living With Hiv In Indonesia: A Qualitative Study. *Int J Environ Res Public Health*. 2022 Jun 4;19(11):6879.
33. Ahmed S, Chase LE, Wagnild J, Akhter N, Sturridge S, Clarke A et al. Community Health Workers And Health Equity In Low- And Middle-Income Countries: Systematic Review And Recommendations For Policy And Practice. *Int J Equity Health*. 2022 Apr 11;21(1):49.
34. Fauk NK, Ward PR, Hawke K, Mwanri L. HIV Stigma And Discrimination: Perspectives And Personal Experiences Of Healthcare Providers In Yogyakarta And Belu, Indonesia. *Front Med (Lausanne)*. 2021 May 12;8.
35. Crocker BCS, Pit SW, Hansen V, John-Leader F, Wright M. A Positive Approach To Adolescent Sexual Health Promotion: A Qualitative Evaluation Of Key Stakeholder Perceptions Of The Australian Positive Adolescent Sexual Health (Pash) Conference. *BMC Public Health*. 2019 Dec 3;19(1):681.
36. Arias-Colmenero T, Pérez-Morente MA, Ramos-Morcillo AJ, Capilla-Díaz C, Ruzafa-Martínez M, Hueso-Montoro C. Experiences And Attitudes Of People With Hiv/Aids: A Systematic Review Of Qualitative Studies. *Int J Environ Res Public Health*. 2020 Jan 19;17(2):639.
37. Akyirem S, Salifu Y, Bayuo J, Duodu PA, Bossman IF, Abboah-Offei M. An Integrative Review Of The Use Of The Concept Of Reassurance In Clinical Practice. *Nurs Open*. 2022 May 11;9(3):1515–35.
38. Bogart Lm, Kgotlaetsile K, Phaladze N, Mosepele M. HIV Self-Testing May Overcome Stigma And Other Barriers To HIV Testing Among Higher-Socioeconomic Status Men In Botswana: A Qualitative Study. *African Journal Of Aids Research*. 2021 Oct 2;20(4):297–306.
39. Smith DK, Henny KD, Weidle PJ. The Evidence Base for Initial Intervention Strategies for Ending the HIV Epidemic in the U.S. *Am J Prev Med*. 2021 Nov;61(5): S1–5.

